


Grass-clover and biogas fertilizers as a climate efficient nutrient supply

Post-treatment of digestate reduce emissions



Nitrogen efficiency*) of different fractions

Spring barley (after injection of manures)			
Digestate	Liquid fraction	NS-solution	Dried fiber
78 %	79 %	98 %	-16 %
Winter wheat (surface-banding)			
44 %	60 %	88 %	

*) relative to mineral fertilizer N

Anaerobic digestion and separation reduced methane emissions from storage, and overall carbon footprint

Kg CO ₂ eq/100 kg grain DM			
Digestate	Liquid fraction	NS-solution	Cattle slurry
17,1	8,5	24,3**)	44,4

***) High field emissions of N₂O due to high amount of liquid.

- Clover grass as a co-substrate in biogas production is an important source of biogenic nitrogen contributing to carbon sequestration and biodiversity
- Anaerobic digestion has a high potential for reduction of GHG emissions
- High nitrogen efficiency of ammonia-rich fractions
- Dried fiber fraction immobilizes nitrogen but is usable as phosphorus source for legume crops

Department of Agroecology

Peter Sørensen
ps@agro.au.dk,
Mobile: +45 2512 5632

Xiaoyi Meng and Søren O. Pedersen
sop@agro.au.dk
Mobile: +45 2812 4304

Marie Trydeman Knudsen
mariet.knudsen@agro.au.dk,
Mobile: +45 2225 8387

Department of Biological and Chemical Engineering

Henrik B. Møller
henrikb.moller@bce.au.dk
Mobile: +45 4053 1596

Innovation Centre for Organic Farming

Erik Fog
eikf@icoel.dk
Mobile: +45 5180 8669